

T H E  
W E S T E R N J O U R N A L

of

MEDICINE AND SURGERY

Edited by

Daniel Drake, M.D.

and

Lunsford P. Yandell, M.D.,

Professors in the Louisville Medical Institute

and

Thomas W. Colescott, M.D.

---

NEW SERIES

VOL. III

---



T H E  
W E S T E R N      J O U R N A L  
OF  
M E D I C I N E   A N D   S U R G E R Y

MARCH, 1845

ART. I.--Observations on the Botany of Illinois, more especially in reference to the Autumnal Flora of the Prairies.  
In a letter to Daniel Drake, M.D., &c.

My Dear Sir:

The interest which you have long taken in every thing relating to the Natural History of the Western States, and the desire which you have so frequently expressed to connect the study of Botany with the pursuit of medicine, will, I trust, be a sufficient apology for my addressing these observations to you. Having moreover, very recently yourself traveled over a part of the country to which they refer, you will be the better enabled to judge of their correctness.

In a tour which I took through the State of Illinois, a few years previous to your own, I had the pleasure of being ac-

*This copy, except for omission of a long-footnote, is page by page and line by line as in the original. Spelling is unchanged.*

Digitized by the Internet Archive  
in 2019 with funding from  
University of Illinois Urbana-Champaign

4 Je 35 Krogh  
accompanied by my brother, and one or two other individuals, who took a considerable interest in the objects which mainly prompted my journey; and traveling in a light covered wagon, well prepared for making extensive collections, and vigilantly on the look-out for every object of interest, I may safely say that few such escaped our observation. Our visit to this interesting region was made in the latter part of summer, and extended so late into the fall, that severe frosts had put an end to vegetation before our return; so that my remarks must be considered as referring to the autumnal Flora of the Prairies, and may not be applicable to that of the spring, or early summer. We entered the State of Illinois from Terre Haute on the Wabash river, near the line dividing that State from Indiana; thence we traveled in nearly a north-western direction to Peoria, on the Illinois river; through Paris, Urbana, Bloomington, and Mackinaw: and returning we took a more southern route through Tremont, Springfield, Hillsborough, Maysville, and Lawrenceville to Vincennes; where we recrossed the Wabash, which here forms the boundary between the States of Indiana and Illinois. This trace extending over a distance of nearly 400 miles, led us through the central portion of the State in two different lines, at a considerable distance apart, and gave us an opportunity of seeing and examining the fact of the country and its productions under a great variety of aspects.

21 Mar 35 q. A. L. Vestal  
In a Geographical point of view, the surface of Illinois may be very appropriately, as it is naturally, divided into three districts. First--The heavily timbered tracts which for the most part occupy the southern portion of the State, bordering on the Ohio river, and which, extending into the middle and northern portions, are found in detached bodies surrounded by prairies, and in these situations are called 'Groves.' These groves are, for the most part, contiguous to, and often bounded by water-courses, which have preserved them from the action of fire. Secondly--The open prairies, of from one to twenty miles in diameter, entirely destitute of



trees, and indeed of all other woody plants, except along the margin of water courses which occasionally pervade them. Thirdly--'The Barrens,' or tracts somewhat intermediate between the two former, being sparsely covered with oak trees of several different kinds, and of considerable size, with a dense undergrowth of various shrubs and annual plants. This third region bears a close relationship, both in appearance and productions, to those districts in Kentucky, which are called 'Barrens'--tracts of country which seem to be in a state of transition from more open prairies to densely timbered forests.

The vegetation of these three districts is, of course, essentially different; but apart from the presence or absence of trees, which constitute the grand feature of distinction, the annual and suffruticose plants are widely different, and indeed in many respects entirely dissimilar. It is, however, to the productions of the open prairies that I shall chiefly confine myself in this communication: and even they vary greatly as the surface of the prairie may be high, rolling, rich and dry, or low, flat, wet and clayey.

The first sight of a prairie with which we were greeted was in the neighborhood of Terre Haute, on the eastern side of the Wabash, and consequently in the State of Indiana. In approaching this new and apparently thriving town, from the east, over the national road, the eye is filled with the prospect of an extensive plain entirely destitute of all timber-trees, and stretching to a great distance both above and below the town. Such a view, agreeable at all times, was peculiarly so as it opened suddenly upon us just after emerging from the heavily wooded forest through which we had traveled all day. The Terre Haute prairie, however, has been all reclaimed, or rather, botanically speaking, desecrated by the hand of man, and no portion of it now remains in a state of nature. Corn, grass, small grain, and other cultivated crops now occupy the hundreds of acres, which lately bloomed and blossomed with indigenous productions; and almost the only relics of these to be seen, were occasion-

...and raised of all other woody ...  
the margin of water courses and ...  
Thirdly--'The ...  
between the two ...  
trees of several ...  
a dense undergrowth of various ...  
This third region bears a close relationship ...  
ance and production, to those districts in ...  
are called 'S ...  
a state of transition from ...  
bered forests.

The vegetation of these three districts ...  
tially different; but apart from the presence ...  
trees, which constitute the ...  
annual and sub-tropical ...  
in many respects ...  
productions of the open ...  
myself in this communication ...  
the surface of the ...  
or low, flat, wet and ...

The first sight of a ...  
was in the neighborhood of ...  
side of the ...  
and below the town. ...  
was peculiarly so as it ...  
emerging from the heavily ...  
had traveled all day. ...  
has been all ...  
created by the ...  
in a state of ...  
cultivated crops now ...  
lately plowed and ...  
and almost the only ...



ally on the road-side, or in fence-corners, a few plants of Verbena stricta and Vernonia corymbosa.

Twenty miles west of the Wabash at this point, we met with the first prairie in a state of nature; and from this, extending northward to the Lakes, and westward to the Mississippi, they continue, increasing in magnitude, and interrupted only by occasional groves of timber, so as to occupy by far the largest portion of the central, eastern, western, and northern portions of the State of Illinois.

On fairly entering the prairie region, and reaching the centre of one of these immense natural meadows, the view presented to the eye of a novice in such scenery, is one of the most pleasing sort. But beautiful, imposing, and even grand as is this spectacle, I must own, that in a botanical point of view, I was disappointed! The Flora of the prairies--the theme of so much admiration to those who view them with an ordinary eye,--does not, when closely examined by the Botanist, present that deep interest and attraction which he has been led to expect. Its leading feature is rather the unbounded profusion with which a few species occur in certain localities, than the mixed variety of many different species occurring any where. Thus from some elevated position in a large prairie the eye takes in at one glance thousands of acres, literally empurpled with the flowering spikes of several species of Liatris, among which the most predominant are L. spicata, L. squarrosa, L. scariosa, L. cylindracea, and L. pycnostachya. In other situations, where a depressed or flattened surface and clayey soil favor the continuance of moisture, a few species of yellow-flowered Coreopsis occur in such profuse abundance as to tinge the entire surface with a golden burnish. The species of this genus more commonly met with in such situations, were Coreopsis trichosperma, C. senifolia, C. tripteris, C. palmata, &c., &c. This peculiarity of an aggregation of individuals of one or more species, to something like an exclusive monopoly of certain localities, obtains even in regard to those plants which are the rarest and least frequently met with; for whenever one specimen was found

ely on the road-side, or in fence-corners,  
Verbes stricta and Verbes stricta

Twenty miles west of the town of ...  
with the first ... in a ...  
tending ... to the ... and ...  
slope, they ... in ...  
only by occasional ... of the ...  
the largest portion of the ...  
ern portions of the ... of Illinois.

On ... entering the ...  
the ... of ...  
sented to the ... of a ...  
most pleasing ... The ...  
as is this ... I was ...  
view, I was ...  
theme of so much ...  
an ordinary eye, ...  
Botanist, present ...  
has been led to ...  
bounded ... with ...  
localities, that the ...  
occurring any where. ...  
large ... the ...  
literally ...  
cies of ...  
spicata, L. ..., L. ...  
nostachya. In other ...  
ed surface and ...  
a few species of ...  
false abundance as to ...  
burnish. The ...  
in such situation, were ...  
O. ..., O. ..., O. ...  
aggregation of ...  
thing like an ...  
even in regard to ...  
frequently not ...

there generally occurred many more in the same immediate neighborhood. The Dalea alopecuroides, (Willd.), which I met with but once, was found in that locality in the greatest abundance. The Satureja hortensis, which I believe is not regarded as indigenous to North America, was seen once by us in the greatest profusion, and that, too, in a situation the least favorable to the idea of its having been introduced--the centre of a large prairie, where no settlement could have been made. At some places between Peoria and Springfield the road-sides and even the beaten path, were so completely covered over with the little Boebera chrysanthemoides, that, trodden under our horses' feet, it exhaled a strong and nauseating odor. In many such localities this noisome weed seems to take the place of the Anthemis cotula and A. arvensis (May-weed and Dog-fennel,) in the more settled portions of the Western States. In the neighborhood of Springfield, again, and especially in the out-lots of that town, we found the ground covered, to the exclusion of almost every other vegetation, with a small species of Ambrosia (A. bidentata) which, at the season in which we saw it, being out of flower, and ripening its dark-colored seed, gave to the common an aspect as dreary as "the bleak and blasted heath where Macbeth met the witches." In illustration of this peculiarity of the Botany of the prairies, I will only further remark that we did not observe the little Erigeron divaricatum until we reached Bloomington, in the commons of which town it is extremely abundant; and that it ceases to occur, or is but rarely seen, a few miles south of that.

There are, indeed, comparatively speaking, but few plants, except the grasses, (which are gregarious every where, and are intermixed in greater or less degree and variety among all the other plants of the prairies,) which may be considered as indigenes of the prairie region generally.--Among these we may mention, as occurring most constantly, and under greater diversity of soil and situation than any others, the Silphium gummiferum, Parthenium integrifolium, Kuhnia critonia, Ceanothus intermedius, (which here takes the



place of C. Americanus in the Barrens of Kentucky,) Prenanthes Illinoensis, Eryngium aquaticum, Petalostemum violaceum, Dracocephalum Virginianum, Baptisia leucantha, several species of Liatris, Coreopsis, Aster,<sup>1</sup> Solidago,<sup>2</sup> Rudbeckia,<sup>3</sup> Helianthus,<sup>4</sup> Pycnanthemum, Gerardia, Pedicularis, Gentiana, &c., &c. Those two beautiful plants, for our knowledge of both of which, I believe, we are indebted to Mr. Nuttall, the Aster sericeus, and Amorpha canescens, are very generally diffused, but not in the same abundance with many others. Indeed, they constitute an exception to the habit of congregation which obtains among so many of their associates.

As is the case, I believe, with the American Flora throughout the United States, and indeed, the whole Continent, the autumnal botany of the prairies exhibits a large preponderance of the Compositae. Besides those already mentioned, we may here enumerate, as of frequent occurrence, Chrysopsis mariana, Helenium autumnale, Boltonia glastifolia and B. asteroides, Bidens frondosa and B. chrysanthemoides, Eupatorium serotinum, E. aromaticum, E. ageratoides, E. purpureum, E. perfoliatum, &c., Cnicus glutinosus, C. Virginianus, C. muticus, C. altissimus, &c., Silphium laciniatum, S. integrifolium, S. terebinthinaceum, &c., Prenanthes aspera, P. virgata, P. racemosa, P. serpentaria, &c., Vernonia fasciculata, V. corymbosa, and one or two other species.

In a farmer's, or rather a grazier's estimation, the grasses would be regarded as the most valuable of the natural productions of the prairies; and we will next mention some of

---

<sup>1</sup>Aster laevis, A. Novae Angliae, A. rigidus, A. gracilis, A. phlogifolius, A. concolor, A. azureus, A. undulatus, A. multiflorus, A. oblongifolius, A. turbinellus, A. carneus, &c.

<sup>2</sup>Solidago rigida, S. nemoralis, S. graminifolia, S. Ridellii, S. serotina, S. speciosa, S. Ohioensis, S. neglecta, &c.

<sup>3</sup>Rudbeckia purpurea, R. laciniata, R. hirta, R. subtomentosa, R. pinnata, &c.

<sup>4</sup>Helianthus angustifolius, H. rigidus, H. occidentalis, H. grosse-serratus, H. tomentosus, H. mollis, H. pubescens, H. microcephalus, H. tomentosus, H. laetiflorus, &c.

place of C. Americanus in the ...  
Antes Illinoisensis, ...  
Dreoscephalus Virginianus, ...  
cles of ...  
thus, ...  
So. These two ...  
of which, I believe, ...  
fer ...  
lused, but not in the ...  
deed, they ...  
tion which ...

As in the case, I believe, ...  
out the United States, ...  
entirely ...  
ance of the ...  
may have ...  
Mariana, ...  
teroides, ...  
serotinus, ...  
F. ...  
muticus, ...  
lum, ...  
P. ...  
corvina, ...

In a ...  
would ...  
function of ...

1. ...  
2. ...  
3. ...  
4. ...  
5. ...  
6. ...  
7. ...  
8. ...  
9. ...  
10. ...

those which are of most frequent occurrence, omitting all reference to the allied tribe of Cyperaceae, but few of which were observed, in consequence, perhaps, of the late season at which our visit was made. Among the most predominant of the Gramineae, on the rich, dry, and rolling prairies are several species of Andropogon, as A. furcatum, A. ciliatum, A. mutans, A. scoparium, &c., Aristida tuberculosa, A. stricta, A. gracilis, &c., Elymus Canadensis, (var. glaucifolius.) E. Virginicus, E. mollis, &c., Trichodium laxiflorum, and Vilfa heterolepsis. In flat and marshy situations these give place to various species of Panicum, as P. geniculatum, P. agrostoides, P. dichotomum, P. virgatum, P. latifolium, and the universally diffused P. crus-galli, Leersia Virginica, and L. oryzoides, Spartina polystachya and S. cynosuroides. All these grasses in their young and tender states are eagerly devoured by cattle: as they become harder and less succulent by age, the coarser are rejected and the more tender are sought for. Among these, I believe, the Vilfa, before mentioned, is a general favorite, both for grazing and for hay. All of them, however, are cut promiscuously for this purpose, and when they occur, as frequently they do, in large natural meadows, occupying the ground to the almost entire exclusion of other vegetables, they yield a productive return to the labor of the mower; and when well cured make excellent hay. Our horses, which had never before been accustomed to any other than the cultivated grasses, ate this natural hay with great avidity. The quality of these grasses, both for pasturage and mowing, is much improved by the burning of the prairies during the winter, which, destroying the dead and dry stems, affords a better and earlier bite in the spring, as well as a cleaner swath for the scythe: and by protecting certain portions of the prairie from the action of fire until the spring or early summer, vegetation is then so much retarded by a 'late burn,' as the settlers call it, as to afford good pasturage throughout the latter part of the season.

To this action of the fires, which probably for ages have annually passed over these plains, consuming in their progress

[illegible]



all relics of vegetable matter, both woody and herbaceous, is perhaps to be mainly ascribed the color of the soil, which for the most part is literally as black as coal, and in some situations of two or three feet in depth. And to this excess of carbonaceous matter, imparted to the soil of these prairies, is it perhaps to be ascribed that their productions, both in cultivated crops and natural growths, are by no means so rank or luxuriant as one might be led to expect. The Indian corn, though well-eared, was not so tall as I have frequently seen it in Kentucky and Ohio, on lands apparently much inferior in fertility; the different kinds of small grain, though heavily-headed, had a much shorter straw; and many of the natural productions, common to the Illinois prairies and the barrens of Kentucky, were less luxuriant in growth than I have observed them to be in the latter district, though the soil of the barrens has not the same appearance of fertility. This subject deserves particular investigation, and an accurate analysis of the prairie soil might lead to very useful practical deductions. One of our fellow-travelers, a farmer by profession, ascribed the appearance, above mentioned, to a 'sourness' in the soil. But the amount of carbonaceous and alkaline matters resulting from such frequent burnings would rather lead to an opposite conclusion.

Among the oeconomical and medicinal plants of the prairies may be mentioned Gentiana ochroleuca, the roots of which have somewhat the bitterness of the officinal species, (G. lutea, of Europe,) Prenanthes serpentaria, several species of Liatris, the tuberous roots of which are possessed of acrid pungent qualities, and Eryngium aquaticum; all these plants have a considerable reputation, which perhaps is but little deserved, against the bites of poisonous serpents, and hence they are known indifferently by the names of 'snake-root,' 'button snake-root,' 'rattle-snake's masterpiece,' &c. Frasera verticillata is not so frequently seen in the more open prairies as in the thinly-wooded barrens. Polygala Senega and Asclepias tuberosa are abundant in both these localities. The different species of Silphium mentioned, exude from their

all kinds of... perhaps to... the most part... tions of... corporation... it seems to be... vated... luxuriant... through... it in... in... heated... provisions... of Kentucky... served... barren... just... sis of the... tutions... sion, a... ness in the... line... rather...

Among the... may be... have some... inter... al... constant... have a... served... they are... 'better... sets... series... and... The...

stems a pearly resinous matter, very similar in appearance and sensible properties to turpentine, and used for the same purposes. The roots of the beautiful Petalostemum violaceum have a warm pungent quality, which suggested its employment, among the thousand other articles, in the treatment of cholera, and the plant is now known on the prairies as the 'cholera-weed.' Our two most valuable indigenous bitters Eupatorium perfoliatum and Sabbatia angularis are abundant, and Aristolochia serpentaria is seen occasionally in the groves, where various species of dogwood (Cornus) are also of frequent occurrence. Mr. J. A. Lapham, of Wisconsin, informs me that in that territory, the Amorpha canescens is called 'lead-plant,' from the circumstance of its growth being considered indicative of the presence of that mineral. If the same sign should hold good in Illinois, the whole of the prairies may one day become a mining region.

Ferns are remarkably rare on the prairies; indeed I do not recollect having met with a single specimen of any species of that extensive tribe in the more open prairies. This may, perhaps, be owing to the absence of that shade and constant moisture in which most of these plants delight. On the skirts of the timbered tracts, several kinds occur, which are usually found in the barrens, as Pteris aquilina, Polypodium dryopteris, and P. hexagonopterum; and in the 'groves' I observed many other species common in the Western States. The same remarks will apply, in a good degree, to the tribe of mosses.

I deem it improper to close these desultory remarks, without giving a catalogue, at least, of other common plants, which presented themselves at different places on our route through the prairies. Some of them may have been already incidentally mentioned, but the most of them occurred under circumstances not calling for particular note or comment. They are given as I find them in my note-book, without any kind of order or arrangement.--



Verbena stricta,  
V. hastata,  
Gerardia purpurea,  
G. flava,  
G. erecta,  
G. auriculata,  
G. quercifolia,  
Petalostemum candidum,  
Desmodium, } of various species.  
Lespedeza, }  
Euphorbia corollata,  
Gaura angustifolia,  
Typha latifolia,  
Cassia chamaecrista,  
C. marilandica,  
Monarda fistulosa,  
Leptandra Virginica,  
Lythrum hyssopifolium,  
Pedicularis pallida,  
Gillenia stipulacea,

Parnassia palustris,  
Gentiana rubricaulis,  
G. quinqueflora,  
Sium latifolium,  
Archemora rigida,  
Artemisia caudata,  
Polygala verticillata,  
P. ambigua,  
P. incarnata,  
Linum rigidum,  
Potentilla fruticosa,  
Psoralea floribunda,  
Boottia sylvestris,  
Plantago cordata,  
P. aristata,  
Cissus Canadensis,  
Chelone glabra,  
Angelica triquinata,  
Epilobium lincaire,  
Lysimachia revoluta, &c.

Doubtless many other species came under our observation, but being so common in other parts of the Western country, I omitted to note them.

In relation to the botany of the prairies, I have only to add a few remarks on the shrubs which are found among them; for although in the more open districts of this kind no ligneous or perennial stems are permitted to escape the ravages of the annual fires which sweep over them, yet on the margins of 'sloughs,' and along the courses of the small streams which occasionally meander through them, clumps of bushes and clusters of shrubbery are always to be found. These 'roughs,' as they are called, furnish welcome retreats to grazing cattle, and sometimes to the traveler's horse, from

1. General  
 2. General  
 3. General  
 4. General  
 5. General  
 6. General  
 7. General  
 8. General  
 9. General  
 10. General  
 11. General  
 12. General  
 13. General  
 14. General  
 15. General  
 16. General  
 17. General  
 18. General  
 19. General  
 20. General  
 21. General  
 22. General  
 23. General  
 24. General  
 25. General  
 26. General  
 27. General  
 28. General  
 29. General  
 30. General  
 31. General  
 32. General  
 33. General  
 34. General  
 35. General  
 36. General  
 37. General  
 38. General  
 39. General  
 40. General  
 41. General  
 42. General  
 43. General  
 44. General  
 45. General  
 46. General  
 47. General  
 48. General  
 49. General  
 50. General  
 51. General  
 52. General  
 53. General  
 54. General  
 55. General  
 56. General  
 57. General  
 58. General  
 59. General  
 60. General  
 61. General  
 62. General  
 63. General  
 64. General  
 65. General  
 66. General  
 67. General  
 68. General  
 69. General  
 70. General  
 71. General  
 72. General  
 73. General  
 74. General  
 75. General  
 76. General  
 77. General  
 78. General  
 79. General  
 80. General  
 81. General  
 82. General  
 83. General  
 84. General  
 85. General  
 86. General  
 87. General  
 88. General  
 89. General  
 90. General  
 91. General  
 92. General  
 93. General  
 94. General  
 95. General  
 96. General  
 97. General  
 98. General  
 99. General  
 100. General

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

[illegible]

that annoying pest of these regions--the prairie fly.\* In these thickets the more common productions are the hazle, (Corylus Americana), three species of sumach, (Rhus glabrum, R. copalinum, and R. aromaticum), several dwarf kinds of plumb, (Prunus), of which the species were not ascertained, two or three varieties of dogwood, (Cornus sericea, C. asperifolia, C. alba, &c.), several species of undetermined willows, (Salix.) Besides these, may be mentioned the Amorpha fruticosa, Zanthoxylum fraxineum, (prickly ash,) Prinos verticillata, Ilex prinoides, Aronia melanocarpa, Spiraea tomentosa and S. salicifolia, Symphorea racemosa, Cephalanthus occidentalis, Rubus

\* Long foot note omitted.

villosus, (blackberry,) Ribes rotundifolium, called Illinois gooseberry, of which the fruit, though spinous, makes a delicious tart; together with various species of wild roses, grape-vines, &c.

Though not properly falling within the compass of this communication, the object of which has been to give some account of the autumnal botany of the prairies, yet before I close it, I will venture to add a few remarks on the forest trees of Illinois. These, in the main, do not differ from the productions of similar districts in the timbered lands of Indiana, Ohio, Kentucky, and Tennessee. In Illinois, the richest groves, interspersed through the prairies, are constituted mainly of the same kind of trees which indicate the best soils generally in the Western States, as black walnut, hickories, hackberry, (Celtis crassifolia), sugar-maple, pawpaw, (Porcelia triloba), &c. The thinner lands are clothed chiefly with oaks of various species, hickories and gums, (Liquidambar

(Footnote continued from Page 195 omitted.)

Page 1 of 1

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020

10/10/2020



styraciflua and Nyssa of two or three species,) whilst the poorest soils, those especially of the 'bushy barrens' and 'oak openings,' are occupied mostly with the different kinds of oak, among which the post-oak, (Quercus obtusiloba,) and black-jack, (Q. ferruginea,) are most prominent. I am able, indeed, to indicate but two trees which are in any way peculiar to the forests of Illinois; and these are the paccan and catalpa. Of these the paccan (Carya olivaeformis,) is found abundantly on the southern borders of the State, where about Shawneetown and other points on the Ohio river, it constituted a large portion of the original forest; and from these districts great quantities of the nuts have been exported. They are not considered, however, to be equal, either in size or flavor, to the paccan-nuts of Texas. The other tree--the catalpa, (Catalpa cordifolia,) I have the authority of General Harrison for saying, is found occasionally, and of large size, in the alluvions of the Wabash river, where he considered it to be certainly a native; in opposition to the opinion of the Abbé Correa, who thought it more probable that the seeds may have been derived from trees planted by the early French settlers of Vincennes and other posts. I have seen this tree in similar alluvions among the dense forests of Henderson county, Kentucky.

Whilst walking over the prairies adjoining the town of Bloomington, in company with our friend Dr. John F. Henry, who resides there, he pointed out to us an extraordinary phenomenon in connexion with vegetation, and one only visible, I suppose, in a prairie country. It was a semicircular, or rather horse-shoe-shaped line of herbage, distinguishable very plainly from the surrounding and included growths, by its darker or deeper green hue. He said that these circles or segments of circles, usually of fifteen or twenty feet diameter, were frequently to be seen in summer, and that it was generally believed they were occasioned by lightning. He described the thunder-storms of this region as sublimely majestic and terrific. We had no opportunity of witnessing this display of Heaven's artillery, during our journey; but

1. The first of these is the fact that the
 2. Government has not been able to
 3. obtain the necessary information
 4. to enable it to make a
 5. proper assessment of the
 6. situation in the country.
 7. This is due to the fact that
 8. the Government has not been
 9. able to establish a
 10. reliable system of
 11. communication with the
 12. various parts of the
 13. country.
 14. The second of these is the fact
 15. that the Government has not
 16. been able to obtain the
 17. necessary funds to
 18. carry out its
 19. policy.
 20. The third of these is the fact
 21. that the Government has not
 22. been able to obtain the
 23. necessary personnel to
 24. carry out its
 25. policy.

1011  
 1012  
 1013  
 1014  
 1015  
 1016  
 1017  
 1018  
 1019  
 1020  
 1021  
 1022  
 1023  
 1024  
 1025  
 1026  
 1027  
 1028  
 1029  
 1030  
 1031  
 1032  
 1033  
 1034  
 1035  
 1036  
 1037  
 1038  
 1039  
 1040  
 1041  
 1042  
 1043  
 1044  
 1045  
 1046  
 1047  
 1048  
 1049  
 1050  
 1051  
 1052  
 1053  
 1054  
 1055  
 1056  
 1057  
 1058  
 1059  
 1060  
 1061  
 1062  
 1063  
 1064  
 1065  
 1066  
 1067  
 1068  
 1069  
 1070  
 1071  
 1072  
 1073  
 1074  
 1075  
 1076  
 1077  
 1078  
 1079  
 1080  
 1081  
 1082  
 1083  
 1084  
 1085  
 1086  
 1087  
 1088  
 1089  
 1090  
 1091  
 1092  
 1093  
 1094  
 1095  
 1096  
 1097  
 1098  
 1099  
 1100  
 1101  
 1102  
 1103  
 1104  
 1105  
 1106  
 1107  
 1108  
 1109  
 1110  
 1111  
 1112  
 1113  
 1114  
 1115  
 1116  
 1117  
 1118  
 1119  
 1120  
 1121  
 1122  
 1123  
 1124  
 1125  
 1126  
 1127  
 1128  
 1129  
 1130  
 1131  
 1132  
 1133  
 1134  
 1135  
 1136  
 1137  
 1138  
 1139  
 1140  
 1141  
 1142  
 1143  
 1144  
 1145  
 1146  
 1147  
 1148  
 1149  
 1150  
 1151  
 1152  
 1153  
 1154  
 1155  
 1156  
 1157  
 1158  
 1159  
 1160  
 1161  
 1162  
 1163  
 1164  
 1165  
 1166  
 1167  
 1168  
 1169  
 1170  
 1171  
 1172  
 1173  
 1174  
 1175  
 1176  
 1177  
 1178  
 1179  
 1180  
 1181  
 1182  
 1183  
 1184  
 1185  
 1186  
 1187  
 1188  
 1189  
 1190  
 1191  
 1192  
 1193  
 1194  
 1195  
 1196  
 1197  
 1198  
 1199  
 1200  
 1201  
 1202  
 1203  
 1204  
 1205  
 1206  
 1207  
 1208  
 1209  
 1210  
 1211  
 1212  
 1213  
 1214  
 1215  
 1216  
 1217  
 1218  
 1219  
 1220  
 1221  
 1222  
 1223  
 1224  
 1225  
 1226  
 1227  
 1228  
 1229  
 1230  
 1231  
 1232  
 1233  
 1234  
 1235  
 1236  
 1237  
 1238  
 1239  
 1240  
 1241  
 1242  
 1243  
 1244  
 1245  
 1246  
 1247  
 1248  
 1249  
 1250  
 1251  
 1252  
 1253  
 1254  
 1255  
 1256  
 1257  
 1258  
 1259  
 1260  
 1261  
 1262  
 1263  
 1264  
 1265  
 1266  
 1267  
 1268  
 1269  
 1270  
 1271  
 1272  
 1273  
 1274  
 1275  
 1276  
 1277  
 1278  
 1279  
 1280  
 1281  
 1282  
 1283  
 1284  
 1285  
 1286  
 1287  
 1288  
 1289  
 1290  
 1291  
 1292  
 1293  
 1294  
 1295  
 1296  
 1297  
 1298  
 1299  
 1300  
 1301  
 1302  
 1303  
 1304  
 1305  
 1306  
 1307  
 1308  
 1309  
 1310  
 1311  
 1312  
 1313  
 1314  
 1315  
 1316  
 1317  
 1318  
 1319  
 1320  
 1321  
 1322  
 1323  
 1324  
 1325  
 1326  
 1327  
 1328  
 1329  
 1330  
 1331  
 1332  
 1333  
 1334  
 1335  
 1336  
 1337  
 1338  
 1339  
 1340  
 1341  
 1342  
 1343  
 1344  
 1345  
 1346  
 1347  
 1348  
 1349  
 1350  
 1351  
 1352  
 1353  
 1354  
 1355  
 1356  
 1357  
 1358  
 1359  
 1360  
 1361  
 1362  
 1363  
 1364  
 1365  
 1366  
 1367  
 1368  
 1369  
 1370  
 1371  
 1372  
 1373  
 1374  
 1375  
 1376  
 1377  
 1378  
 1379  
 1380  
 1381  
 1382  
 1383  
 1384  
 1385  
 1386  
 1387  
 1388  
 1389  
 1390  
 1391  
 1392  
 1393  
 1394  
 1395  
 1396  
 1397  
 1398  
 1399  
 1400  
 1401  
 1402  
 1403  
 1404  
 1405  
 1406  
 1407  
 1408  
 1409  
 1410  
 1411  
 1412  
 1413  
 1414  
 1415  
 1416  
 1417  
 1418  
 1419  
 1420  
 1421  
 1422  
 1423  
 1424  
 1425  
 1426  
 1427  
 1428  
 1429  
 1430  
 1431  
 1432  
 1433  
 1434  
 1435  
 1436  
 1437  
 1438  
 1439  
 1440  
 1441  
 1442  
 1443  
 1444  
 1445  
 1446  
 1447  
 1448  
 1449  
 1450  
 1451  
 1452  
 1453  
 1454  
 1455  
 1456  
 1457  
 1458  
 1459  
 1460  
 1461  
 1462  
 1463  
 1464  
 1465

in two or three instances afterwards, I think, we observed this singular appearance of the grass on the prairies, indicating what might, perhaps, without impiety, be called 'the foot-prints of the Deity!'

Very respectfully and truly, I am,

My dear Sir, yours,

C. W. SHORT

Professor Drake,  
Medical Institute of Louisville, }

February, 1845.

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

[illegible]

• 3 •